

Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Organochlorine Compounds and Ultrasound Measurements of Fetal Growth in the INMA Cohort (Spain)

Maria-Jose Lopez-Espinosa, Mario Murcia, Carmen Iñiguez, Esther Vizcaino, Olga Costa, Ana Fernández-Somoano, Mikel Basterrechea, Aitana Lertxundi, Mònica Guxens, Mireia Gascon, Fernando Goñi-Irigoyen, Joan O. Grimalt, Adonina Tardón, and Ferran Ballester

Table of Contents

Detailed modeling procedure of mixed-effect models and growth curves

Rationale

General model description

Conditional and unconditional centiles

Steps of the modeling procedure

Table S1. Summary of the models by cohort. The INMA Project, 2003-2008 (Spain).

Figure S1. Fetal growth curves for estimated fetal weight (EFW) in the four INMA-cohorts, 2003-2008 (Spain)

Figure S2. Fetal growth curves for abdominal circumference (AC) in the four INMA-cohorts, 2003-2008 (Spain).

Figure S3. Fetal growth curves for biparietal diameter (BPD) in the four INMA-cohorts, 2003-2008 (Spain)

Figure S4. Fetal growth curves for femur length (FL) in the four INMA-cohorts, 2003-2008 (Spain). There were not different FL curves by sex since sex did not enter in any model.

Details on the multiple imputation (MI) modeling

Imputation method

Software and statistical packages

Number of imputed datasets and iterations

Variables included in the imputation procedure

Heterogeneity in the imputation modeling

Criteria of inclusion

Diagnostics

Table S2. Number (%) of imputed values and regression model for each variable. The INMA Project, 2003-2008 (Spain).

Table S3. Pearson's correlations between OCs in maternal and umbilical cord serum. The INMA Project, 2003-2008 (Spain).

Figure S5. Sensitivity analyses of the associations between OC concentrations and fetal growth measurements between 0-12 weeks of gestation. The INMA Project, 2003-2008 (Spain).

Figure S6. Sensitivity analyses of the associations between OC concentrations and fetal growth measurements between 12-20 weeks of gestation. The INMA Project, 2003-2008 (Spain).

Figure S7. Sensitivity analyses of the associations between OC concentrations and fetal growth measurements between 20-34 weeks of gestation. The INMA Project, 2003-2008 (Spain).

Footnote of the Supplemental Material, Figures S5-S7

References